

**POULTRY SULFA - sulfamethazine sodium, sulfamerazine sodium and sulfaquinoxaline sodium powder, for solution**  
Alpharma Inc. Animal Health

**SULFONAMIDE**

**For Oral Veterinary Use Only**

**For Use in Drinking Water Only**

As an aid in the control of coccidiosis and acute fowl cholera in chickens and acute fowl cholera and coccidiosis in turkeys, when caused by pathogens susceptible to sulfamethazine, sulfamerazine and sulfaquinoxaline.

**ANTIMICROBIAL SOLUBLE POWDER**

**For Use in Chickens and Turkeys**

**THIS PACKET CONTAINS:**

78 grams Sodium Sulfamethazine Activity

78 grams Sodium Sulfamerazine Activity

39 grams Sodium Sulfaquinoxaline Activity

**CAUTION**

Federal law prohibits the extralabel use of this product in lactating dairy cattle.

**WARNING (HUMAN FOOD)**

Do not treat chickens or turkeys within 14 days of slaughter for food. Do not medicate chickens or turkeys producing eggs for human consumption.

**DIRECTIONS FOR USE**

	PROPORTIONER SOLUTION (1 oz/gal)	TANK MIX
.04% Solution	Add one pack to 1 gallon (3.8 liters)	Add one pack to 128 gallons
.025% Solution	Add one pack to 1.6 gallons (6.1 liters)	Add one pack to 206 gallons

**MAKE FRESH SOLUTION DAILY.** If improvement is not noted in 72 hours, consult a poultry veterinarian or poultry diagnostic laboratory for diagnosis. During treatment use only medicated water unless otherwise directed. For control of disease outbreaks medication should be initiated as soon as diagnosis is determined. Treated animals must actually consume enough medicated water to provide a therapeutic dose. Do not mix or administer in galvanized containers. Dispose of any waste or unused portions properly.

**PRECAUTION**

May cause toxic reactions unless drug is evenly mixed in water at dosages indicated and used according to label directions.

**DIRECTIONS**

**Acute Fowl Cholera - TURKEYS AND CHICKENS**

As an aid in the control of acute fowl cholera caused by *Pasteurella multocida* susceptible to sulfamethazine, sulfamerazine and sulfaquinoxaline. Provide medicated water (.04% solution) for 2-3 days. If disease recurs, repeat treatment.

**Coccidiosis - TURKEYS**

As an aid in the control of coccidiosis caused by *Eimeria meleagritidis* and *E. adenoides* susceptible to sulfamethazine, sulfamerazine and sulfaquinoxaline. Provide medicated water (.025% solution) for 2 days, then plain water for 3 days, then medicated water (.025% solution) for 2 days, then plain water for 3 days, then medicated water (.025% solution) for 2 days. Repeat if necessary. DO NOT CHANGE LITTER.

**Coccidiosis - CHICKENS**

As an aid in the control of coccidiosis caused by *Eimeria tenella* and *E. necatrix* susceptible to sulfamethazine, sulfamerazine and sulfaquinoxaline. Provide medicated water (.04% solution) for 2-3 days, then plain water for 3 days, then medicated water (.025% solution) for 2 days. If bloody droppings appear, repeat at .025% level for 2 more days. DO NOT CHANGE LITTER.

**Keep Out of Reach of Children**



Alpharma Inc.

Bridgewater, New Jersey 08807

Restricted Drug (California) - Use only as directed.

NADA 100-094, Approved by FDA

Trademarks registered by Alpharma Inc.  
9001BF 0911

**PRINCIPAL DISPLAY PANEL - 195 G PACKET**  
**RUSSELL®**

**PoultrySulfa<sup>®</sup>**  
**Sulfonamide**

**For Oral Veterinary Use Only**  
**For Use in Drinking Water Only**

As an aid in the control of coccidiosis and acute fowl cholera in chickens and acute fowl cholera and coccidiosis in turkeys, when caused by pathogens susceptible to sulfamethazine, sulfamerazine and sulfaquinoxaline.

# ANTIMICROBIAL SOLUBLE POWDER

## For Use in Chickens and Turkeys

THIS PACKET CONTAINS:

## 78 grams Sodium Sulfamethazine Activity

### 78 grams Sodium Sulfamerazine Activity

### 39 grams Sodium Sulfaquinolaxline Activity

**CAUTION:** Federal law prohibits the extralabel use of this product in lactating dairy cattle.

**Keep Out of Reach of Children**

## Take Time

## Observe Label

### Directions

Alpharma Inc.

Bridgewater, New Jersey 08807


Restricted Drug (California) - Use only as directed.

NADA 100-094, Approved by FDA

Trademarks registered by Alpharma Inc.

9001BF 0911

Directions for use:		
<b>Warning</b> <b>(Human Food)</b> Do not feed chickens or turkeys within 14 days of slaughter for human consumption. Do not feed to other animals (chickens or turkeys) producing eggs for human consumption.	<b>1. PROPIONATORNER SOLUTION (1 ounce)</b> <b>1.4% Solution:</b> Add one pump to 1.4 gallons (1.1 liter).	<b>TANK MIX</b> Add one pump to 128 gallons.
	<b>2.5% Solution:</b> Add one pump to 2.5 gallons (1.9 liter).	Add one pump to 256 gallons.
	<b>MAKE FRESH SOLUTION DAILY:</b> If propionate is not noted in 72 hours, contact a poultry veterinarian or poultry diagnostic laboratory for diagnosis. During treatment use only medicated water unless otherwise indicated. Do not use medicated water if the diagnosis should be initiated as soon as diagnosis is determined. Treat animal until actually healthy. Do not use medicated water to provide a therapeutic or other benefit in a generalized condition. Discontinue if any signs or untoward properties appear.	
<b>PRECAUTIONS:</b> May cause toxic reactions unless drug is evenly mixed in water. All drugs indicated and used according to label directions.		



**RUSSELL & RUSSELL**

# PoultrySulfa™

**Sulfamonomide**  
**For Chick Veterinary Use Only**  
**Use in Drinking Water Only**  
 As an aid in the control of  
 coccidiosis and acute foot  
 chills in chickens and adult foot  
 chills in turkeys, when  
 caused by pathogens susceptible to  
 sulfamonomide, sulfamonomide and  
 sulfapropylenol.

**ANTIBACTERIAL  
 SOLUBLE POWDER**


**Use in Chickens and Turkeys**

**THIS PACKET CONTAINS:**  
 75 grams Sulfamonomide Sulfazine Activity  
 75 grams Sulfamonomide Sulfazine Activity  
 75 grams Sulfamonomide Sulfazine Activity  
 36 grams Sulfamonomide Sulfazine Activity

**Keep Out of Reach of Children**

**Take One**  
**Observe Label**  
**Directions**

**CAUTION:** Federal law prohibits the  
 withdrawal of this product in feeding  
 areas.



**ALPHARMA**  
 ALPHARMA INC.  
 Kalamazoo, MI 49001

**Restained Dog (California):** Use only as directed.  
**NADA 105-084, Approved for use in:**  
 California, Nevada, Arizona, New Mexico, Utah, Idaho, Oregon, Washington, Alaska, Hawaii, Puerto Rico, U.S. Possessions, U.S. Virgin Islands, Guam, American Samoa, Northern Mariana Islands, and the Federated States of Micronesia.

**PRINCIPAL DISPLAY PANEL - 40 PACK CARTON**  
**RUSSELL®**

## Sulfonamide

## Antimicrobial

## For Oral Veterinary

**Use Only**

## PoultrySulfa®

Follow directions for use and mixing instructions on packet label.

**CAUTION:** For use in drinking water only. Keep out of reach of children.

Store in a cool, dry place.

**Contents: 40 packs,**

each packet contains:

78 grams Sodium Sulfamethazine Activity

78 grams Sodium Sulfamerazine Activity

39 grams Sodium Sulfaquinoxaline Activity

**Lot:**

**Exp.:**

**Take Time**

**Observe Label**

**Directions**

Restricted Drug (California) - Use only as directed.

9001CL 0911

NADA #100-094, Approved by FDA

Trademarks registered by Alpharma Inc.

Alpharma Inc.

Bridgewater, New Jersey 08807



Sulfonamide  
Antimicrobial  
For Oral Veterinary  
Use Only

# PoultrySulfa®

Follow directions for use and mixing instructions on packet label.

**CAUTION:** For use in drinking water only. Keep out of reach of children.  
Store in a cool, dry place.

**Lot:**

**Exp.:**



Alpharma Inc.  
Bridgewater, New Jersey 08807

**Contents: 40 packs,**  
each packet contains:

78 grams Sodium Sulfamethazine Activity

78 grams Sodium Sulfamerazine Activity

39 grams Sodium Sulfaquinoxaline Activity

**Take Time**



**Observe Label  
Directions**

Restricted Drug (California) - Use only as directed.

9001CL 0911

NADA #100-094, Approved by FDA

Trademarks registered by Alpharma Inc.